



Calf STARTER Plus

(FOR RUMINANTS ONLY)



NUTRIENT ANALYSIS

Protein, not less than	15%
Fat, not less than	6%
Fiber, not more than	1%
Calcium (min)8%
Calcium (max).....	1.2%
Phosphorous, not less than.....	1%
Potassium, not less than.....	5.6%
Cobalt, not less than.....	4.1 ppm
Copper, not less than.....	19 ppm
Manganese, not less than.....	215 ppm
Selenium, not less than.....	4.5 ppm
Zinc, not less than.....	254 ppm
Iodine, not less than.....	9 ppm
Vitamin A, not less than.....	200,000 IU/lb
Vitamin D3, not less than.....	20,000 IU/lb
Vitamin E, not less than.....	80 IU/lb
Bio-Mos.....	Added

INGREDIENTS

Beet molasses, Concentrated separator by-products, Corn distillers grain, Navy bean by-products, Hydrolyzed vegetable fat (preserved with Ethoxyquin), Monocalcium-calcium phosphate, Calcium carbonate, Magnesium oxide, Zinc Sulfate, Zinc Oxide, Zinc polysaccharide complex, Copper Sulfate, Copper polysaccharide complex, Ferrous sulfate, Magnanous oxide, Maganese sulfate, Ethylenediamine dihydroiodide, Cobalt sulfate, Sodium selenite, Vitamin A acetate, Vitamin D3 supplement, aspergillus oryzae and saccharomyces cervaise fermentation culture (Progro 45), Flax seed, Diatomaceous Earth, Bio-Mos (mannan oligosaccharides)



FEEDING DIRECTIONS

Feed one Calf hutch bucket per hutch. Feed one 20 lb tub to 1-5 calves. Feed one 60 lb tub to 1-5 calves. Place the tubs where cattle tend to congregate. Tubs should be fed continuously along with additional forages and water. If additional mineral or vitamins are desired offer a loose mineral free choice.

IMPORTANT

Progro 45 works as a catalyst which dramatically increases the numbers of bacteria by breaking down the complex carbohydrates so that bacteria can feed on the simple sugars the enzymes produce.

INTRODUCTION PERIOD

During the introduction period tubs should be placed approximately 50-75 feet near the area that cattle congregate. Move tubs to accommodate consumption, if consumption is less than expected, move tubs nearer to the area that the cattle congregate. If consumption is more than expected move tubs further from congregational area. Additional feed may be necessary if consumption becomes excessive. DO NOT PROVIDE TUBS TO CATTLE OF POOR CONDITION WITHOUT PROVIDING SUPPLEMENTAL FEED. IF CONSUMPTION IS EXCESSIVE FOR MORE THAN TWO WEEKS REEVALUATE THE TOTAL FEEDING PROCESS.

Caution: DO NOT FEED CALF STARTER PLUS TO SHEEP (OR RELATED SPECIES), PRODUCT CONTAINS COPPER SULFATE WHICH IS TOXIC TO SHEEP.

Manufactured by:



149 RR Ave | Wyndmere, ND 58081 | www.tublicks.com

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed.

_____ Product Code # 9071-15	15 lbs (6.8 kg) Net Weight or Bulk
_____ Product Code # 9071-20	20 lbs (9.1 kg) Net Weight or Bulk
_____ Product Code # 9071-60	60 lbs (27.2 kg) Net Weight or Bulk

Lot No. _____