



# Calf Weaner

(FOR RUMINANTS ONLY)



## NUTRIENT ANALYSIS

|                                 |               |
|---------------------------------|---------------|
| Protein, not less than .....    | 15%           |
| Fat, not less than .....        | 6%            |
| Fiber, not more than .....      | 1%            |
| Calcium (min) .....             | 8%            |
| Calcium (max).....              | 1.2%          |
| Phosphorous, not less than..... | 1%            |
| Potassium, not less than.....   | 5.6%          |
| Cobalt, not less than.....      | 4.1 ppm       |
| Copper, not less than.....      | 19 ppm        |
| Manganese, not less than.....   | 215 ppm       |
| Selenium, not less than.....    | 4.5 ppm       |
| Zinc, not less than.....        | 254 ppm       |
| Iodine, not less than.....      | 9 ppm         |
| Vitamin A, not less than.....   | 200,000 IU/lb |
| Vitamin D3, not less than.....  | 20,000 IU/lb  |
| Vitamin E, not less than.....   | 80 IU/lb      |
| Bio-Mos.....                    | Added         |

## INGREDIENTS

Beet molasses, Concentrated separator by-products. Corn distillers grain. Navy bean by-products. Hydrolyzed vegetable fat (preserved with Ethoxyquin). Monocalcium-calcium phosphate. Calcium carbonate. Magnesium oxide. Zinc sulfate. Zinc oxide. Zinc polysaccharide complex, Copper Sulfate, Copper polysaccharide complex. Ferrous sulfate. Magnanous oxide. Maganese sulfate. Ethylenediamine dihydroiodide. Cobalt sulfate. Sodium selenite. Vitamin A acetate, Vitamin D3 supplement. Vitamin E supplement, concentrated aspergillus oryzae, saccharomyces cervasiae fermentation culture, Urea, Diatomaceous Earth (anti-caking agent) added, Bio-mos (mannan oligosaccharides), Flax seed.



## FEEDING DIRECTIONS

Feed one tub per 20 head of cattle. Additional tubs may be needed when there are smaller calves with larger cattle. Place the tubs where cattle tend to congregate. Tubs should be fed continuously along with additional forages and water. If additional mineral or vitamins are desired, offer a loose mineral free choice

### IMPORTANT

Consumption of Calf Weaner will vary depending on quality feed and availability of other feeds. Cattle conditions and weather conditions will also affect consumption.

## INTRODUCTION PERIOD

During the introduction period, tubs should be placed approximately 50 to 75 feet near the area that cattle congregate. Move tubs to accommodate consumption, if consumption is less than expected, move tubs nearer to the area that cattle congregate. If consumption is more than expected move tubs further from congregational area. Additional feed may be necessary if consumption becomes excessive. DO NOT PROVIDE TUBS TO CATTLE OF POOR CONDITION WITHOUT PROVIDING SUPPLEMENTAL FEED. IF CONSUMPTION IS EXCESSIVE FOR MORE THAN TWO WEEKS REEVALUATE THE TOTAL FEEDING PROGRAM.

**Caution: DO NOT FEED TUB LICKS TO SHEEP (OR RELATED SPECIES), CALF WEANER CONTAINS COPPER SULFATE WHICH IS TOXIC TO SHEEP.**

Manufactured by:



149 RR Ave | Wyndmere, ND 58081 | www.tublicks.com

*This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed.*

Product Code # 9073-250 \_\_\_\_\_

Product Code # 9073-500 \_\_\_\_\_

Lot No. \_\_\_\_\_

250 Lbs (113.4 kg), 500 Lbs (226.8 kg) Net Weight or Bulk