



DAIRY- LICKS 19% AN

(FOR RUMINANTS ONLY)

NUTRIENT ANALYSIS

| | |
|-------------------------------|-----------------------|
| Crude protein, minimum..... | 19.00% |
| Crude fat, minimum..... | 5.00% |
| Crude fiber, maximum..... | 2.00% |
| Calcium (Ca), minimum..... | 2.00% |
| Calcium (Ca), maximum..... | 3.00% |
| Phosphorous (P), minimum..... | .10% |
| Magnesium (Mg), minimum..... | .80% |
| Potassium (K), minimum..... | 6.25% |
| Sulfur (S), minimum..... | .45% |
| Zinc (Zn), minimum..... | 2000 ppm |
| Copper (Cu), minimum..... | 500 ppm |
| Manganese (Mn), minimum..... | 1500 ppm |
| Cobalt (Co), minimum..... | 40 ppm |
| Iodine (I), minimum..... | 50 ppm |
| Selenium (Se), minimum..... | 6 ppm |
| Vitamin A, minimum..... | 63,000 Int. Units/Lb. |
| Vitamin D3, minimum..... | 35,000 Int. Units/Lb. |
| Vitamin E, minimum..... | 2,000 Int. Units/Lb. |

INGREDIENTS

Beet molasses, Concentrated separator by-products. Corn distillers grain. Navy bean by-products. Hydrolyzed vegetable fat (preserved with Ethoxyquin). Monocalcium-calcium phosphate. Calcium carbonate. Magnesium oxide. Zinc sulfate. Zinc oxide. Zinc polysaccharide complex, Copper Sulfate, Copper polysaccharide complex. Ferrous sulfate. Magnanous oxide. Maganese sulfate. Ethylenediamine dihydroiodide. Cobalt sulfate. Sodium selenite. Vitamin A acetate, Vitamin D3 supplement. Vitamin E supplement, concentrated aspergillus oryzae, saccharomyces cervasiae fermentation culture). Diatomaceous Earth (anti-caking agent) added.



FEEDING DIRECTIONS

DAIRY-LICKS 19% AN Nutritional supplement is designed to be fed at the start of the dry period, along with a long stem, low calcium forage. Continue feeding until freshening. Place containers in dry spots near loafing areas and watering locations. Feed at the rate of one container for every 20-30 head, with a minimum of two (2) containers in each lot or pasture to ensure proper accessibility. Free-choice consumption will depend on the size of the animals fed, seasonal weather conditions, as well as quality, source and availability of other feeds. Cattle typically consume between 0.5 and 1.0 pounds per head daily.

IMPORTANT

ALL CONSUMPTION FIGURES HERE ARE APPROXIMATE. PROVIDE FREE ACCESS TO CLEAN, FRESH WATER AT ALL TIMES. DAIRY-LICKS 19% AN NUTRITIONAL SUPPLEMENT CONTAINS NO SALT. PROVIDE SALT FREE CHOICE.

INTRODUCTION PERIOD

During the introduction period, tubs should be placed approximately 50 to 75 feet near the area that cattle congregate. Move tubs to accommodate consumption, if consumption is less than expected, move tubs nearer to the area that cattle congregate. If consumption is more than expected move tubs further from congregational area. Additional feed may be necessary if consumption becomes excessive. DO NOT PROVIDE TUBS TO CATTLE OF POOR CONDITION WITHOUT PROVIDING SUPPLEMENTAL FEED. IF CONSUMPTION IS EXCESSIVE FOR MORE THAN TWO WEEKS REEVALUATE THE TOTAL FEEDING PROGRAM.

Caution: DO NOT FEED DAIRY-LICKS 19% AN TO SHEEP (OR RELATED SPECIES), DAIRY-LICKS 19% AN CONTAINS COPPER SULFATE WHICH IS TOXIC TO SHEEP.

Manufactured by:



149 RR Ave | Wyndmere, ND 58081 | www.tublicks.com

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed.

Product Code # 8070

Lot No. _____

150 Lbs (69.0 kg), 250 Lbs (113.4 kg), 500 Lbs (226.8 kg) Net Weight or Bulk