



Whinney- Licks 15%

(FOR EQUINE)

NUTRIENT ANALYSIS

Crude protein, minimum.....	15.00%
Crude fat, minimum.....	5.00%
Crude fiber, maximum.....	1.50%
Calcium (Ca), minimum	1.00%
Phosphorous (P), minimum	1.00%
Magnesium (Mg), minimum	1.75%
Potassium (K), minimum.....	6.00%
Sulfur (S), minimum45%
Zinc (Zn), minimum.....	900 ppm
Copper (Cu), minimum.....	250 ppm
Manganese (Mn), minimum	850 ppm
Cobalt (Co), minimum.....	3 ppm
Iodine (I), minimum	9 ppm
Selenium (Se), minimum.....	11 ppm
Vitamin A, minimum	36,000 Int. Units/Lb.
Vitamin D3, minimum.....	4,000 Int. Units/Lb.
Vitamin E, minimum.....	245 Int. Units/Lb.
Biotin, minimum.....	2.5 mg/Lb.
Glucosamine Hydrochloride.....	170 mg/Lb

INGREDIENTS

Beet molasses, Concentrated separator by-product, Flaxseed, Corn distillers grains, Hydrolyzed vegetable fat (preserved with Ethoxyquin), Monocalcium-dicalcium phosphate, Calcium Carbonate, Magnesium oxide, Vitamin A acetate, D-activated animal sterol (source of Vitamin D3 activity), Vitamin E supplement, Menadoine demethyl pyrimidinal bisulphate (source of Vitamin K activity), Niacin supplement, Calcium pantothenate, Riboflavin supplement, Vitamin B-12 supplement, Folic acid, Pyridoxine hydrochloride, Thiamine mononitrate, Biotin, Zinc sulfate, Zinc oxide, Copper sulfate, Ferrous sulfate, Manganous oxide, Ethylenediamine digyroidide, Cobalt carbonate, Sodium selenite, Diatomaceous Earth (anti-caking agent) added, concentrated aspergillus oryzae, saccharomyces cervasiae fermentation culture),. Glucosamine Hydrochloride.



FEEDING DIRECTIONS

A supplemental feed for horses. Place tubs where horses tend to gather. Feed continuously along with additional forages or to horses on pasture.

IMPORTANT

Consumption of Whinney Licks will vary depending on quality feed and availability of other feeds. Condition of horses and weather conditions will also affect consumption.

INTRODUCTION PERIOD

During the introduction period, tubs should be placed approximately 50 to 75 feet near the area that horses congregate. Move tubs to accommodate consumption, if consumption is less than expected, move tubs nearer to the area that horses congregate. If consumption is more than expected move tubs further from congregational area. Additional feed may be necessary if consumption becomes excessive. DO NOT PROVIDE TUBS TO HORSES OF POOR CONDITION WITHOUT PROVIDING SUPPLEMENTAL FEED. IF CONSUMPTION IS EXCESSIVE FOR MORE THAN TWO WEEKS REEVALUATE THE TOTAL FEEDING PROGRAM.

Caution: DO NOT FEED WHINNEY-LICKS TO SHEEP (OR RELATED SPECIES), WHINNEY-LICKS CONTAINS COPPER SULFATE WHICH IS TOXIC TO SHEEP.

Manufactured by:



149 RR Ave | Wyndmere, ND 58081 | www.tublicks.com

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed.

_____ Product Code # 7010-020	20 lbs (9.1 kg) Net Weight or Bulk
_____ Product Code # 7010-060	60 lbs (27.2 kg) Net Weight or Bulk
_____ Product Code # 7010-150	150 lbs (69.0 kg) Net Weight or Bulk
_____ Product Code # 7010-250	250 lbs (113.4 kg) Net Weight or Bulk

Lot No. _____